

The Potential for Stem Cells in Cerebral Palsy – Piecing Together the Puzzle

*Stuart. D. Faulkner (PhD)¹²³, *Crystal A. Ruff (PhD)¹²³, Michael G. Fehlings (MD. PhD. FRCSC, FACS)¹²³⁴

¹Division of Genetics and Development, Toronto Western Research Institute, Toronto, Ontario, Canada

²Institute of Medical Science, University of Toronto, Toronto, Ontario, Canada

³Spinal Program, University Health Network, Toronto Western Hospital, Toronto, Ontario, Canada

⁴Division of Neurosurgery, University of Toronto, Toronto, Ontario, Canada

* Authors contributed equally to this manuscript

Correspondence should be addressed to:

Michael G. Fehlings MD. PhD. FRCSC. FACS

Division of Neurosurgery

University Health Network

Toronto Western Hospital

399 Bathurst St. 4WW-449

Toronto, ON, M5T 2S8, Canada

Tel: (416) 603-5229

Fax: (416) 603-5745

E-mail: michael.fehlings@uhn.on.ca

1) Summary

The substantial socio-economic burden of a diagnosis of Cerebral Palsy (CP), coupled with a positive anecdotal and media spin on stem cell treatments, drives many affected families to seek information and treatment outside of the current clinical and scientific realm. Preclinical studies using several types of stem and adult cells – including mesenchymal stem cells (MSCs), neural precursor cells (NPCs), olfactory ensheathing glia (OEG) and Schwann cells (SCs) - have demonstrated some regenerative and functional efficacy in neurological paradigms. This paper describes the most common cell types investigated for transplant *in vivo* and summarizes the current state of early phase clinical trials. It investigates the most relevant and promising co-administered therapies – including rehabilitation, drug targeting, magnetic stimulation and bioengineering approaches. We highlight the need for adjunctive combinatorial strategies to successfully transfer stem cell treatments from bench to bedside.

2) Introduction

Cerebral Palsy (CP) comprises a heterogeneous group of non-progressive developmental disorders and is marked by loss of neuromotor function. This is often accompanied by demyelination of cerebral white matter tracts and loss of axons and grey matter. Children with CP often demonstrate concomitant deficits of motor control, cognition, learning and other complex neurological functions. Stem cell therapy has become a topic of interest in the popular media and stem cell transplantation is often perceived as a “cure” for CP and other neurological conditions. This misperception (along with anecdotal claims of efficacy and denial of negative side effects by unregulated foreign clinics purporting to offer “stem cell therapy”) presents an unrealistic bias, influencing stakeholders’ perceptions of availability, efficacy and safety of “stem cell treatments” for CP (see review¹). Despite promise in preclinical and clinical trials using stem cells for CP, there remains a knowledge gap surrounding the optimal source and type of cells, timing of treatment and possible mechanisms of action. We present here an updated summary of the current state of stem cell science for CP and describe current relevant clinical trials. This article highlights the need for additional optimization before stem cell treatments can be fully realised as a therapeutic option for CP.

I. The Etiology of CP

CP is a heterogeneous disability and its etiology is equally multivariate. It remains poorly understood despite a large body of research and literature on the topic. The current “multi-hit” hypothesis surrounding its etiology maintains that alignment of multiple probabilities, rather than a single risk factor, is responsible for CP onset. Indeed, several neonates survive major insults without any evidence of impairment, while others develop severe infarcts almost spontaneously. While an exhaustive description of all causative mechanisms for CP is beyond the scope of this manuscript, the most common risk factors are:

- Preterm birth (and associated complications)
- Intrauterine Growth Restriction
- Maternal infection - viral, bacterial and protozoan (most commonly TORCHS)

T – Toxoplasmosis/ Toxoplasma Gondii

O– “Other” infections (Coxsackievirus, Varicella-Zoster Virus, HIV and Parvovirus B19)

R – Rubella

C – Cytomegalovirus

H – Herpes Simplex Virus

S - Syphilis

- Perinatal and Intrapartum difficulties (ex. hypoxic/ischemic injury)
- Other congenital infections (ex. sepsis, meningitis, encephalitis, tetanus and chorioamnionitis^{2–5})

II. CP Pathophysiology

Inflammation, which may occur through transplacental cytokine passage or by fetal production⁶, plays a key role in the pathophysiology of CP. Increased serum concentrations of B-lymphocyte chemo-attractant, ciliary neurotrophic factor (CNTF), epidermal growth factors, IL-12, IL-15, monocyte chemo-attractant protein-3, and others were found to be present in children with CP⁷, and there is evidence for maturation-dependent changes in both fetal brain structure and response to immune challenge⁸. Pro inflammatory cytokines - which may act 1) directly on pre-oligodendrocytes, 2) via secondary, effector-mediated processes, or most likely, 3) through a combination of the two - have been linked to periventricular leukomalacia (PVL), the most common physiological cause of CP⁹.

Due to GluR subunit composition, the immature oligodendrocyte is intrinsically vulnerable to ischemic and inflammatory insult, particularly to Glutamate-mediated excitotoxicity^{10, 11}. Oligodendrocytes express GluR3 and GluR4 receptors during development, but – unlike pre-progenitors or mature oligodendrocytes - fail to express GluR2. This absence of GluR2 leads to enhanced Ca²⁺ membrane permeability, increased Ca²⁺ influx and resultant excitotoxic cell death. Therefore, as a result, after global ischemic hypoperfusion, periventricular pre-oligodendrocytes, which are most distal to major cerebral vasculature, are selectively abolished, causing focal necrotic lesions and later widespread apoptosis associated with PVL¹². These areas spatially correspond with descending corticospinal tracts, leading to the neuromotor symptoms in

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284> the most common CP phenotype. Likewise, unilateral hemorrhagic or hypoxic/ischemic

injury leads to neuronal cell death in motor control areas, resulting in the second most common subtype of CP, spastic hemiplegia (Figure 1).




Figure 1: Schematic Diagram Representing the Pathophysiology of Hemiplegic and Diplegic Cerebral Palsy

A) Hypoxic Stroke causes hemispheric loss of cortical tissue perfused by the occluded artery. Affected regions overlap the corticospinal tracts, which control lower limbs, trunk and forelimbs. Resulting gross neuromuscular deficits on the contralateral side of the body translate to hemiplegic cerebral palsy. B) Chronic hypoperfusion causes ischemia in developing periventricular watershed regions of the brain. Since the pre-oligodendrocyte is particularly susceptible to ischemic injury, supporting glial cells are preferentially ablated in this region, causing periventricular leukomalacia. These watershed regions overlap with descending corticospinal axons, and without sufficient signal propagation, many motor neurons are damaged. Altogether, these changes lead to spastic diplegia.

3) Current Clinical Treatments for CP

At this time, non-surgical clinical treatments for CP are restricted primarily to rehabilitation and supportive strategies. Many of the current clinical treatments for CP attempt to promote recovery through stimulation of plasticity via activity-dependent changes in excitability and synaptic strength; this enhanced plasticity could be complimentary to regenerative therapeutics. Future regenerative stem cell strategies are likely to be used in combination with these rehabilitation and physiological measures to facilitate clinical translation.

A. Rehabilitation

Rehabilitation therapy is employed under the premise that regularly utilized neuromotor pathways will strengthen circuit plasticity and also that repeatedly used muscle systems will lead to enhanced co-ordination and function. It is currently the most effective treatment for CP symptoms.

B. Physiotherapy and Occupational Therapy

Physiotherapy (PT) and Occupational Therapy (OT) are the primary health care professions associated with rehabilitation. PT encompasses strengthening of ability, co-ordination, function and movement while OT focuses on application of skills toward

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284> activity-defined goals. However, optimal frequency and duration of PT and OT remain to be fully elucidated (reviewed in ¹³).

C. Constraint-Induced Movement Therapy (CIMT) and Bimanual Therapy

CIMT is employed in hemiplegic CP and consists of unilateral constraint on the unaffected side – usually via casting – coupled with shaping techniques and repetitive practice. Alternatively, bimanual training attempts to balance interhemispheric neuromotor competition and involves practiced use of integrated dexterous tasks using both affected and unaffected limbs. Both show promising clinical results, despite little mechanistic preclinical animal data to support or explain their efficacy ^{14–16}.

D. Gaming

Virtual reality simulators and active video games (AVGs) require advanced and coordinated leg, arm and body movement. AVGs can increase a child's participation in active rehabilitation, daily living and engagement in social activities. Furthermore, it can improve self image and mental health. A recent systematic review showed an average increase in energy expenditure of 222% ($\pm 100\%$) during AVG play in typically developing children¹⁷.

E. Intrathecal Baclofen Pumps

Baclofen is a gamma-aminobutyric acid (GABA) agonist, which impedes the release of excitatory neurotransmitters in the spinal cord, and is utilized to treat generalized

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284> spasticity or dystonia. Baclofen can be infused via an intrathecal infusion pump,¹⁸

reducing some side effects such as sedation, confusion, dizziness, ataxia, weakness, nausea, hypotension and parasthesia¹⁹.

F. Botulinum Toxin (Botox)

Botulinum toxin (Botox) is a neurotoxin that causes reversible neuromotor blockage (3-6 months duration). It can provide temporary relief of tone, manage spasticity-related pain, promote longitudinal muscle growth and improve general motor function in targeted areas. Strong safety and efficacy evidence exists for its clinical use and low dose injections at multiple sites can be used safely in children^{20, 21}.

G. Transcranial Magnetic Stimulation (TMS)

TMS utilizes an electromagnetic coil, placed on the scalp, to create electromagnetic pulses that can focally and non-invasively depolarize specific neuronal cortical and sub-cortical targets. Repeated TMS can induce long-lasting activational changes (Long term potentiation (LTP) and long term depression (LTD)) in selected brain areas, particularly those affecting neuromotor control. In animals, intermittent, high-frequency stimulation is generally associated with LTP and longer periods of lower frequency stimulation produce LTD. Although this is promising in model systems, the frequencies and intensities of signals used for LTP can lead to seizures in humans²²; thus, this treatment needs further work to establish an optimal protocol for delivery.

4) Stem Cells

There are many types of mature and stem cells which have been used for experimental (and early-phase clinical) treatments in CP and other CNS paradigms. These studies utilize intraparenchymal, intrathecal and intravenous delivery.

I. Pluripotent Stem Cells

A. Embryonic Stem Cells (ESCs)

Embryonic stem cells (ESCs) are derived from the inner cell mass of the developing blastocyst and are pluripotent in nature. The capacity of ESCs to propagate in culture over many passages provides an almost unlimited supply of cells. However, their allogenicity necessitates immunosuppression in transplant scenarios. Furthermore, their ethical derivation (generally requiring destruction of the embryo) remains highly controversial, making alternative sources more attractive.

B. Induced Pluripotent Stem Cells (iPSCs)

Induced pluripotent stem cells (iPSCs) can be derived from human adult somatic tissue and arguably resemble embryonic stem cells morphologically, antigenically and phenotypically, offering an alternative source of pluripotent cells. By upregulating “Yamanaka” transcription factors OCT4, c-Myc, Sox2 and KLF4,^{23, 24} iPSCs have been generated from several species, including mouse and human. As they can be prepared from an individual’s own cells, iPSCs can potentially reduce the possibility of immune

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

rejection while circumventing ethical issues that arise from cells directly derived from embryonic origin. The most efficient current derivation methods use viral vectors, and are thus unsuitable for human translation. These methods are further hampered by low iPSC yield and the recent discovery that human iPSCs often display somatic coding mutations as well as epigenetic variability²⁵. The recent use of non-integrating vectors, such as the piggybac transposon²⁶, excisable recombinase systems such as Cre/picornavirus systems²⁷, ribonucleic acid²⁸, small molecules²⁹ and non-integrating viruses³⁰, have addressed some of the aforementioned concerns.

C. Epigenetic Variability

Although iPSCs share the same genetic signature as their somatic counterparts, differences with pluripotency are at least partially due to epigenetic variability. Indeed, methylation patterns and high-order chromatin organization in large heterochromatin domains are differentially expressed in pluripotent and differentiated cells³¹. Furthermore, this epigenetic variability is cell-type specific. iPSCs derived from distinct somatic populations reveal similarly distinguishable methylation patterns, characteristics and differentiation potential, despite identical iPSC derivation^{32, 33}. Functionally, synovial tissue MSCs were more effective in cartilage repair than MSCs derived from bone marrow, muscle or fat, indicating that this epigenetic variability in genetically identical specimens can influence performance *in vivo*³⁴. Differential methylation patterns are also observed between ES and iPS-derived pluripotent stem cells³⁵, indicating that

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284> epigenetic, as well as genetic, reprogramming may be necessary for successful human translation.

D. Fate Restriction

Despite their relative ease of expansion, pluripotent cells display a high degree of plasticity and can form teratomas upon transplant. For ESC or iPSC-derived cells to be used in successful cell therapy, they must be differentiated into multipotent or fate-restricted progenitors. Differentiation can be guided by using a number of factors; however, NPCs created via the default pathway seem to be most effective at turning into myelinating oligodendrocytes *in vivo*³⁶. The default pathway differentiates pluripotent cells along a neural lineage by first fate-restricting to a primitive, LIF-dependent, state before reaching a definitive, more differentiated LIF-independent phenotype³⁷.

Alternatively, methods such as direct lineage reprogramming, or “trans-differentiation” have been used to laterally reprogram cells of one mature phenotype into another. Built on analysis of differential transcriptional regulation made popular by iPS technology, trans-differentiation has been used in neural paradigms to derive neurons or tri-potent neural precursors from fibroblasts^{38–41}. Functional neurons have also been generated from hepatocytes, which originate from a different germ layer than neural tissue³⁸. *In vivo*, neuroectodermal trans-differentiation has been explored between post-mitotic callosal and corticofugal neurons³⁹ as well as retinal photoreceptors⁴⁰.

II. Adult Stem Cells

A. Neural Precursor Cells (NPCs)

Neural precursor cells (NPCs) are naturally found in the sub-granular region of the hippocampus, sub-ependymal zone of the spinal cord and the sub ventricular area of adult brain. They can also be derived from fetal and embryonic brain tissue. They can generate all three neural cell types - neurons, astrocytes and oligodendrocytes. In addition, NPCs can be differentiated from pluripotent cell sources via the default pathway³⁷. When transplanted *in vivo* without pre-conditioning, default pathway derived-NPCs almost exclusively differentiate into glial subtype cells, to the exclusion of neurons³⁶. Other groups have used different methods to derive alternative cell types from pluripotent sources (Reviewed in ⁴¹) and the first investigation using iPS-derived NPCs has shown functional efficacy in non-human primate models, albeit with tri-potential *in vivo* differentiation⁴².

Preclinically, transplantation of NPCs or more differentiated glial progenitor cells (GPCs) – both adult and pluripotent-derived - in animal models of injury and dysmyelination is associated with migration of cells to the site of injury, remyelination, functional improvement and low rates of tumorigenesis^{43–47} (Figure 2).




Figure 2: The Function of Neural Precursor Cells for Injury Repair

Hypoxic and hemodynamic insults lead to robust demyelination. B) Transplanted neural precursors can migrate to sites of demyelination and replace lost oligodendrocytes. C) This remyelination often leads to functional recovery and can re-establish axonal signalling. D) Cell transplant also frequently leads to partial restoration of brain tissue.

Furthermore, there is a natural but functionally insufficient propensity for expansion and mobilization of endogenous NPC populations following neurological damage⁴⁸. Although some work has been done to activate or preserve these endogenous populations following injury via drug therapy^{49, 50}, these studies have been unable to successfully

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284> mobilize numbers of cells comparable to that which can be injected from an expanded exogenous population.

The propensity of many NPCs and GPCs to differentiate into glial supporting cells (to the exclusion of neurons) *in vivo* is advantageous in the context of CP, where demyelination predominates and additional neuronal formation or motor neuron connectivity can be deleterious. Glial progenitor cells (GPCs) arising from neural stem cells are arguably an ideal source of safe and pure oligodendrocyte producing cells for remyelination in demyelination disorders such as hereditary leukodystrophies and CP (see ⁵¹ for review). Hence, NPCs (or their derivatives) show the most potential for positive functional results with clinical translation, despite a relatively small number of studies using them in neural injury models.

B. Mesenchymal Stem Cells (MSCs)

Within the bone marrow and umbilical cord exist two major subsets of cells: 1) hematopoietic and endothelial stem cells (CD34+), which form blood cells within the body, and 2) non-hematopoietic mesenchymal stem cells (MSCs, CD34-), which generate bone, cartilage, fat, blood vessels, and connective tissue ^{52, 53}. MSCs are spindle-shaped, fibroblast-like multipotent adult stem cells with limited capacity for self-renewal, which comprise a small population of the adherent stromal cell fraction. MSCs are traditionally derived from bone marrow, umbilical cord or placenta, although they have also been found in synovium, fat, blood vessels and articular cartilage ^{54–58}.

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

Immunogenicity of MSCs is comparatively low; they exhibit minimal expression of Human Leukocyte Antigen (HLA)-B⁵⁹, and can demonstrate immunosuppressive and anti-inflammatory activity. In the transplant microenvironment, MSCs can influence several other cell types- either through cell-to-cell contact, trophic modulation or a combination of both - including T cells, natural killer cells, dendritic cells, monocytes, and neutrophils. MSCs are known to secrete prostaglandins, VEG-F, FGF, indoleamine 2,3-dioxygenase, soluble HLA-G5, IL-6, IL-10, TGB- β 1, HGF, BiNOS, and heme oxygenase- 1 (^{60, 61} reviewed in ⁶²). Although there is some evidence to suggest they can form neural derivatives *in vitro*, this does not occur *in vivo* to a meaningful extent⁶³. Hence, the potential therapeutic benefit of MSCs appears to be mainly via immune mediation, angiogenesis, chemotropism and cellular sparing rather than via trans-differentiation into neural myelinating cells.

In preclinical studies of CP, intraperitoneal, intracardiac and intrahemispheric injection of MSCs has resulted in highly variable outcomes (reviewed in ⁶⁴). Nevertheless, because hematopoietic niche-derived cells have been used for decades in cell transplant paradigms unrelated to CP, such as leukemia and autoimmune disorders⁶⁵, they have promptly moved laterally into safety trials for CP. Considering the inconsistency associated with cell-sourcing and selection conditions currently employed, MSCs, at present, have limited application in chronic cases of human CP.

III. Mature Cells

A. Peripheral Glial Cells

Schwann Cells (SCs) and Olfactory Ensheathing Glia (OEG) are generally agreed to be myelinating cells, found in the PNS and olfactory system respectively. However, they are not “true” stem cells. These cells have myelination capacity and can provide trophic and structural support following injury⁶⁶. Historically, SCs were the cell type first studied in transplant paradigms⁶⁷. Because they create potent trophic and physical substrates for axonal growth, proximal sensory and propriospinal axons readily enter and rarely leave SC grafts^{68, 69}. The olfactory bulb and the lamina propria of the olfactory mucosa are the main sources of OEG. OEG have shown some promise in SCI models, but mixed results in brain injury models. Their limited application in *in vivo* brain injury models might be due to their origin or culture conditions (reviewed in⁷⁰). Consequently, strategies that employ SC transplant generally involve co-transplanted growth factors, biomaterials⁷¹ or cells such as OEGs.

5) Current Clinical Stem Cell Trials for CP

Currently, eight (one suspended, one completed with results) early phase clinical trials are using stem cells for treatment of CP or its antecedents (according to www.clinicaltrials.gov⁷²), using umbilical cord or bone marrow-derived MSCs. Most

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284> consist of treatment in the chronic phase of injury and primarily recruit non gender specific adolescents ~1 to 12 years of age, with diagnosed CP (often mixed etiology), that lack seizures. The only study (from Korea) with results was a double-blinded randomized trial. One hundred and five participants were between 10 months and 10 years of age, with a male: female of ~2:1. They were separated into groups which received either: 1) Allogeneic umbilical cord blood infusion + Erythropoietin (EPO) injection + active rehabilitation, 2) EPO + active rehabilitation, or 3) active rehabilitation, in the chronic phase of injury. Assessment was made at baseline and 1, 3 and 6 months post intervention. Primary outcome measures of motor function and standardized gross motor function were improved in the MSC group compared to other groups. Secondary outcome measures of cognition and neurodevelopmental outcome were also improved in the MSC group compared to other groups (clinical trials identifier: NCT01193660). Despite these results, neurodevelopmental time points beyond 12 months post treatment, coupled with secondary analysis, is required to fully elucidate the potential of this study. Seven clinical trials using NPCs exist in various CNS disorders. The first phase I clinical trial using human ESC derived GPCs for spinal cord injury (clinical trials identifier; NCT01217008) showed promise, but was halted in 2011 due to reported financial difficulties. Since then, fetal-derived NPCs and GPCs have been used as the primary source of NPCs in clinical trials. In other demyelination disorders such as advanced Batten's disease (neuronal ceroid lipofuscinosis) and Pelizaeus-Merzbacher disease, a high intraparenchymal dose of NPCs in multiple injection sites was well tolerated in phase I studies^{73, 74}.

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284> Furthermore, there is a growing subversive culture of “stem cell clinics”, which operate

outside of geographic regulation. These offer “stem cell therapies” for a variety of conditions, based on non-existent or sparse preclinical data. Anecdotal claims of efficacy from such clinics often feed a worrying trend of “Stem Cell Tourism”, with individuals and parents seeking treatment or “cures” for their children with developmental disabilities. While a detailed discussion is elsewhere (⁷⁵), financial costs (often exceeding \$30,000) and individual risks are high.

6) Combinatorial Strategies

The extent of brain injury in CP is often not constrained to just axonal loss and demyelination. Cognitive and behavioural deficits commonly involve grey matter damage; therefore strategies that preserve all three are important to consider in the context of current rehabilitative strategies and future stem cell approaches. Current strategies for CNS repair have largely focused on two separate approaches to promote recovery: exogenous cell transplantation and endogenous cell stimulation. However, it is increasingly likely that additional approaches, which increase plasticity and structural support (bioscaffolds) as well as immunomodulation, will be required in combination with stem cell transplantation.

I. Growth Factors and Engineered Stem Cells

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

Growth factors, such as Neurotrophin-3 (NT-3) (which induces neuronal differentiation), granulocyte colony-stimulating factor (G-CSF), stem cell factor and brain derived neurotrophic factor (BDNF), are associated with improved transplant survival and migration. Furthermore, multifaceted strategies involving NPCs, combined with growth factors (NT-3 or multiple trophic factors), chondroitinase or Schwann cells, have increased functional recovery in spinal cord injury (see review ⁷⁶).

II. Bioscaffolds

Cross linked hyaluronan (HA)-based synthetic extracellular matrix (sECM)⁷⁷⁻⁷⁹ and alginate scaffolds⁷⁶ can secrete growth factors over days and weeks post-implantation. Their capacity to secrete substrate over a sustained period of time ameliorates the need for repeated invasive injections. In the context of brain injury, where extensive loss of cerebral tissue exists, providing a physical substrate to bridge the tissue gap during cellular regeneration strategies is vital. In experimental SCI, self-assembling peptide nanofibers (SAPs) from peptide amphiphile molecules have been used to bridge the spinal cord cavity. Additionally, use of these SAPs resulted in axonal elongation and glial scar inhibition.⁸⁰ Evidence is emerging that SAPs, from 16 peptide (RADA16-1) molecules, can be effective in brain injury models. They have demonstrated integration into the cavity and a reduction in the number of immune reactive cells local to the lesion (see review⁸¹). Multifaceted strategies of polymer scaffolds, growth factors and NPC transplantation in experimental SCI⁷⁶ indicate an important technological advancement for improved structural support and cell survival in clinical applications of brain injury.

III. Clinical Combinatorial Treatments

Although stem cell science shows great promise for reducing CP symptom load, clinical rehabilitative strategies are currently the *only* standardized treatments that provide benefit. Clinical combinatorial treatments are growing in popularity in attempts to see additive or synergistic effects. Studies using functional neuromuscular electrical stimulation in combination with BTXA injection or wrist splints (see ⁸² for review), or direct current stimulation (tDSC) combined with CIMT ⁸³ have shown promise, but require greater numbers for complete evaluation.

While one current clinical trial using combinatorial strategies of EPO and cord blood MSCs in CP claims positive results, it is yet unclear what additional benefits combined treatments may have.

7) Conclusions

Despite preclinical progress, media focus and clinical promise, stem cell therapy for CP remains incomplete and in need of optimization. Easily-expandable, patient-specific cellular therapies are on the horizon, but are currently several years (if not decades) away from clinical reality. Despite this, parents of children with developmental disabilities are often unsure and often misguided as to what treatment options exist and where to seek them. The future of stem cell therapy for CP is evolving. A much better

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

understanding of basic stem cell biology must be established in the field, with appropriate cell type, source, derivation methods and safety profile optimized prior to clinical trials. Also, the correct timing, route of delivery and patient demographic must be taken into consideration when designing early clinical studies. Finally, successful regenerative medicine will involve a multifaceted, combinatorial approach combining current clinical rehabilitation, bioengineering and regenerative stem cell based strategies. Stem cell therapy shows substantial promise in the context of regenerative medicine. Once these important challenges and milestones are overcome, stem cell transplantation for CP could benefit millions of affected children worldwide.

8) References

1. Patoine B: NerveCenter: media focus on “miracle cure” for cerebral palsy pits science vs. hype. *Ann Neurol* 2009; 66:A9–11
2. Wu YW, Colford JM Jr: Chorioamnionitis as a risk factor for cerebral palsy: A meta-analysis. *JAMA* 2000; 284:1417–1424
3. Crichton JU, Dunn HG, McBurney AK, et al.: Long-term effects of neonatal jaundice on brain function in children of low birth weight. *Pediatrics* 1972; 49:656–670
4. Gordon AL, English M, Tumaini Dzombo J, et al.: Neurological and developmental outcome of neonatal jaundice and sepsis in rural Kenya. *Trop Med Int Health* 2005; 10:1114–1120
5. Lawn JE, Cousens S, Zupan J: 4 million neonatal deaths: when? Where? Why? *Lancet* 2005; 365:891–900
6. Burd I, Al B, Chai J, et al.: Yudkoff M, Elovitz MA: Inflammation-induced preterm birth alters neuronal morphology in the mouse fetal brain. *J Neurosci Res* 2010;
7. Zupan V, Gonzalez P, Lacaze-Masmonteil T, et al.: Periventricular Leukomalacia: Risk Factors Revisited. *Developmental Medicine & Child Neurology* 1996; 38:1061–1067[cited 2013 Jan 15]
8. Meyer U, Nyffeler M, Engler A, et al.: The time of prenatal immune challenge determines the specificity of inflammation-mediated brain and behavioral pathology. *J Neurosci* 2006; 26:4752–4762
9. Burd I, Balakrishnan B, Kannan S: Models of Fetal Brain Injury, Intrauterine Inflammation, and Preterm Birth. *American Journal of Reproductive Immunology* 2012; 67:287–294[cited 2012 Nov 1]
10. Fern R, Moller T: Rapid Ischemic Cell Death in Immature Oligodendrocytes: A Fatal Glutamate Release Feedback Loop. *J Neurosci* 2000; 20:34–42[cited 2010 May 24]
11. Itoh T, Beesley J, Itoh A, et al.: AMPA glutamate receptor-mediated calcium signaling is transiently enhanced during development of oligodendrocytes. *J Neurochem* 2002; 81:390–402

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

12. Back SA, Luo NL, Mallinson RA, et al.: Selective vulnerability of preterm white matter to oxidative damage defined by F2-isoprostanes. *Ann Neurol* 2005; 58:108–120[cited 2011 Aug 4]
13. Palisano RJ, Begnoche DM, Chiarello LA, et al.: Amount and focus of physical therapy and occupational therapy for young children with cerebral palsy. *Phys Occup Ther Pediatr* 2012; 32:368–382
14. Gordon AM, Chinnan A, Gill S, et al.: “Both constraint-induced movement therapy and bimanual training lead to improved performance of upper extremity function in children with hemiplegia”. *Developmental Medicine & Child Neurology* 2008; 50:957–958[cited 2012 Dec 13]
15. Gordon AM, Hung Y-C, Brandao M, et al.: Bimanual training and constraint-induced movement therapy in children with hemiplegic cerebral palsy: a randomized trial. *Neurorehabil Neural Repair* 2011; 25:692–702
16. Charles JR, Gordon AM: A repeated course of constraint-induced movement therapy results in further improvement. *Developmental Medicine & Child Neurology* 2007; 49:770–773[cited 2012 Dec 13]
17. Biddiss E, Irwin J: Active video games to promote physical activity in children and youth: A systematic review. *Archives of Pediatrics and Adolescent Medicine* 2010; 164:664–672
18. Dralle D, Müller H, Zierski J, et al.: INTRATHECAL BACLOFEN FOR SPASTICITY. *The Lancet* 1985; 326:1003[cited 2012 Nov 21]
19. Goldstein EM: Spasticity management: an overview. *J Child Neurol* 2001; 16:16–23
20. Bakheit AM, Severa S, Cosgrove A, et al.: Safety profile and efficacy of botulinum toxin A (Dysport) in children with muscle spasticity. *Dev Med Child Neurol* 2001; 43:234–238
21. Kawamura A, Campbell K, Lam-Damji S, et al.: A randomized controlled trial comparing botulinum toxin A dosage in the upper extremity of children with spasticity. *Dev Med Child Neurol* 2007; 49:331–337
22. Wassermann EM: Risk and safety of repetitive transcranial magnetic stimulation: Report and suggested guidelines from the International Workshop on the Safety of Repetitive Transcranial Magnetic Stimulation, June 5-7, 1996. *Electroencephalography and Clinical Neurophysiology - Evoked Potentials* 1998; 108:1–16

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

23. Takahashi K, Yamanaka S: Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* 2006; 126:663–676[cited 2010 May 2]
24. Takahashi K, Tanabe K, Ohnuki M, et al.: Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007; 131:861–872[cited 2010 May 7]
25. Gore A, Li Z, Fung H-L, et al.: Somatic coding mutations in human induced pluripotent stem cells. *Nature* 2011; 471:63–67
26. Woltjen K, Michael IP, Mohseni P, et al.: piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells. *Nature* 2009; 458:766–770[cited 2010 Feb 18]
27. Loh Y-H, Yang JC, De Los Angeles A, et al.: Excision of a viral reprogramming cassette by delivery of synthetic Cre mRNA. *Curr Protoc Stem Cell Biol* 2012; Chapter 4:Unit4A.5
28. Warren L, Manos PD, Ahfeldt T, et al.: Highly Efficient Reprogramming to Pluripotency and Directed Differentiation of Human Cells with Synthetic Modified mRNA. *Cell Stem Cell* 2010; 7:1–13
29. Ichida JK, Blanchard J, Lam K, et al.: A small-molecule inhibitor of tgfbeta signaling replaces sox2 in reprogramming by inducing nanog. *Cell Stem Cell* 2009; 5:491–503[cited 2010 May 9]
30. Fusaki N, Ban H, Nishiyama A, et al.: Efficient induction of transgene-free human pluripotent stem cells using a vector based on Sendai virus, an RNA virus that does not integrate into the host genome. *Proceedings of the Japan Academy, Series B* 2009; 85:348–362
31. Wen B, Wu H, Loh Y-H, et al.: Euchromatin islands in large heterochromatin domains are enriched for CTCF binding and differentially DNA-methylated regions. *BMC Genomics* 2012; 13:566
32. Kim K, Zhao R, Doi A, et al.: Donor cell type can influence the epigenome and differentiation potential of human induced pluripotent stem cells. *Nat Biotechnol* 2011; 29:1117–1119
33. Kim K, Doi A, Wen B, et al.: Epigenetic memory in induced pluripotent stem cells. *Nature* 2010; 467:285–290

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

34. Koga H, Muneta T, Nagase T, et al.: Comparison of mesenchymal tissues-derived stem cells for in vivo chondrogenesis: suitable conditions for cell therapy of cartilage defects in rabbit. *Cell Tissue Res* 2008; 333:207–215
35. Deng J, Shoemaker R, Xie B, et al.: Targeted bisulfite sequencing reveals changes in DNA methylation associated with nuclear reprogramming. *Nat Biotechnol* 2009; 27:353–360[cited 2010 Jun 20]
36. Rowland JW, Lee JJ, Salewski RP, et al.: Generation of Neural Stem Cells from Embryonic Stem Cells Using the Default Mechanism: In Vitro and In Vivo Characterization [Internet]. *Stem Cells and Development* 2011; [cited 2011 Sep 20] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21604925>
37. Smukler SR, Runciman SB, Xu S, et al.: Embryonic stem cells assume a primitive neural stem cell fate in the absence of extrinsic influences. *J Cell Biol* 2006; 172:79–90[cited 2010 May 2]
38. Marro S, Pang ZP, Yang N, et al.: Direct lineage conversion of terminally differentiated hepatocytes to functional neurons. *Cell Stem Cell* 2011; 9:374–382
39. Rouaux C, Arlotta P: Direct lineage reprogramming of post-mitotic callosal neurons into corticofugal neurons in vivo. *Nat Cell Biol* 2013;
40. Montana CL, Kolesnikov AV, Shen SQ, et al.: Reprogramming of adult rod photoreceptors prevents retinal degeneration. *Proc Natl Acad Sci USA* 2013;
41. Ruff CA, Fehlings MG: Neural stem cells in regenerative medicine: bridging the gap. *Panminerva Med* 2010; 52:125–147[cited 2010 Jun 8]
42. Kobayashi Y, Okada Y, Itakura G, et al.: Pre-Evaluated Safe Human iPSC-Derived Neural Stem Cells Promote Functional Recovery after Spinal Cord Injury in Common Marmoset without Tumorigenicity. *PLoS ONE* 2012; 7:e52787
43. Karimi-Abdolrezaee S, Eftekharpour E, Wang J, et al.: Delayed transplantation of adult neural precursor cells promotes remyelination and functional neurological recovery after spinal cord injury. *J Neurosci* 2006; 26:3377–3389[cited 2010 May 10]
44. Karimi-Abdolrezaee S, Eftekharpour E, Wang J, et al.: Synergistic effects of transplanted adult neural stem/progenitor cells, chondroitinase, and growth factors promote functional repair and plasticity of the chronically injured spinal cord. *J Neurosci* 2010; 30:1657–1676[cited 2010 Mar 15]

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

45. Windrem MS, Schanz SJ, Guo M, et al.: Neonatal chimerization with human glial progenitor cells can both remyelinate and rescue the otherwise lethally hypomyelinated shiverer mouse. *Cell Stem Cell* 2008; 2:553–565[cited 2010 Feb 18]
46. Obenaus A, Dilmac N, Tone B, et al.: Long-term magnetic resonance imaging of stem cells in neonatal ischemic injury. *Ann Neurol* 2011; 69:282–291[cited 2011 Jul 4]
47. Fujimoto Y, Abematsu M, Falk A, et al.: Treatment of a Mouse Model of Spinal Cord Injury by Transplantation of Human iPS Cell-derived Long-term Self-renewing Neuroepithelial-like Stem Cells [Internet]. *Stem Cells* (Dayton, Ohio) 2012; [cited 2012 Mar 19] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22419556>
48. Zawadzka M, Rivers LE, Fancy SPJ, et al.: CNS-resident glial progenitor/stem cells produce Schwann cells as well as oligodendrocytes during repair of CNS demyelination. *Cell Stem Cell* 2010; 6:578–590[cited 2010 Oct 14]
49. Hunt J, Cheng A, Hoyle A, et al.: Cyclosporin A has direct effects on adult neural precursor cells. *J Neurosci* 2010; 30:2888–2896
50. Wang J, Gallagher D, DeVito LM, et al.: Metformin activates an atypical PKC-CBP pathway to promote neurogenesis and enhance spatial memory formation. *Cell Stem Cell* 2012; 11:23–35
51. Goldman SA, Nedergaard M, Windrem MS: Glial progenitor cell-based treatment and modeling of neurological disease. *Science* 2012; 338:491–495
52. Erices A, Conget P, Minguell JJ: Mesenchymal progenitor cells in human umbilical cord blood. *Br J Haematol* 2000; 109:235–242
53. Nieda M, Nicol A, Denning-Kendall P, et al.: Endothelial cell precursors are normal components of human umbilical cord blood. *Br J Haematol* 1997; 98:775–777
54. Barbero A, Ploegert S, Heberer M, et al.: Plasticity of clonal populations of dedifferentiated adult human articular chondrocytes. *Arthritis Rheum* 2003; 48:1315–1325[cited 2010 Oct 2]
55. Battula VL, Bareiss PM, Trembl S, et al.: Human placenta and bone marrow derived MSC cultured in serum-free, b-FGF-containing medium express cell surface frizzled-9 and SSEA-4 and give rise to multilineage differentiation. *Differentiation* 2007; 75:279–291[cited 2010 Oct 2]

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

56. De Bari C, Dell'Accio F, Tylzanowski P, et al.: Multipotent mesenchymal stem cells from adult human synovial membrane. *Arthritis Rheum* 2001; 44:1928–1942[cited 2010 Oct 2]
57. Weiss ML, Medicetty S, Bledsoe AR, et al.: Human umbilical cord matrix stem cells: preliminary characterization and effect of transplantation in a rodent model of Parkinson's disease. *Stem Cells* 2006; 24:781–792[cited 2010 Oct 2]
58. Zuk PA, Zhu M, Mizuno H, et al.: Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue Eng* 2001; 7:211–228[cited 2010 Oct 2]
59. Isa A, Nehlin JO, Sabir HJ, et al.: Impaired Cell Surface Expression of HLA-B Antigens on Mesenchymal Stem Cells and Muscle Cell Progenitors [Internet]. *PLoS One* 2010; 5[cited 2013 Jan 14] Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2878340/>
60. Liao W, Xie J, Zhong J, et al.: Therapeutic effect of human umbilical cord multipotent mesenchymal stromal cells in a rat model of stroke. *Transplantation* 2009; 87:350–359[cited 2010 May 11]
61. Chen J, Zhang ZG, Li Y, et al.: Intravenous administration of human bone marrow stromal cells induces angiogenesis in the ischemic boundary zone after stroke in rats. *Circ Res* 2003; 92:692–699[cited 2010 May 11]
62. Prockop DJ: Repair of tissues by adult stem/progenitor cells (MSCs): controversies, myths, and changing paradigms. *Mol Ther* 2009; 17:939–946[cited 2010 Oct 2]
63. Meletis K, Frisé J: Blood on the tracks: a simple twist of fate? *Trends Neurosci* 2003; 26:292–296[cited 2010 May 12]
64. Bennet L, Tan S, Van den Heuvel L, et al.: Cell therapy for neonatal hypoxia–ischemia and cerebral palsy. *Annals of Neurology* 2012; 71:589–600[cited 2013 Jan 15]
65. Ikehara S: Bone marrow transplantation: a new strategy for intractable diseases. *Drugs Today* 2002; 38:103–111
66. Pearse DD, Sanchez AR, Pereira FC, et al.: Transplantation of Schwann cells and/or olfactory ensheathing glia into the contused spinal cord: Survival, migration, axon association, and functional recovery. *Glia* 2007; 55:976–1000[cited 2010 May 7]

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

67. Duncan ID, Aguayo AJ, Bunge RP, et al.: Transplantation of rat Schwann cells grown in tissue culture into the mouse spinal cord. *J Neurol Sci* 1981; 49:241–252
68. Raisman G: Use of Schwann cells to induce repair of adult CNS tracts. *Rev Neurol (Paris)* 1997; 153:521–525[cited 2010 Oct 10]
69. Campbell G, Hutchins K, Winterbottom J, et al.: Upregulation of activating transcription factor 3 (ATF3) by intrinsic CNS neurons regenerating axons into peripheral nerve grafts. *Exp Neurol* 2005; 192:340–7
70. Mackay-Sim A, St John JA: Olfactory ensheathing cells from the nose: clinical application in human spinal cord injuries. *Exp Neurol* 2011; 229:174–180
71. Chen G, Hu Y, Wan H, et al.: Functional recovery following traumatic spinal cord injury mediated by a unique polymer scaffold seeded with neural stem cells and Schwann cells. *Chin Med J* 2010; 123:2424–2431
72. Home - ClinicalTrials.gov [Internet]. [cited 2013 Jan 15] Available from: <http://www.clinicaltrials.gov/>
73. StemCells, Inc. : Groundbreaking Science. Breakthrough Medicine. [Internet]. [cited 2013 Jan 17] Available from: <http://www.stemcellsinc.com/>
74. StemCells finds early data from spinal cord injury trial promising - San Francisco Business Times [Internet]. [cited 2012 Sep 20] Available from: <http://www.bizjournals.com/sanfrancisco/blog/biotech/2012/09/stemcells-neural-stem-cells-spinal-cord.html>
75. Lau D, Ogbogu U, Taylor B, et al.: Stem Cell Clinics Online: The Direct-to-Consumer Portrayal of Stem Cell Medicine. *Cell Stem Cell* 2008; 3:591–594
76. Hwang DH, Kim HM, Kang YM, et al.: Combination of multifaceted strategies to maximize the therapeutic benefits of neural stem cell transplantation for spinal cord repair. *Cell Transplant* 2011; 20:1361–1379
77. Prestwich GD: Hyaluronic acid-based clinical biomaterials derived for cell and molecule delivery in regenerative medicine. *J Control Release* 2011; 155:193–199
78. Wang Y, Wei YT, Zu ZH, et al.: Combination of hyaluronic acid hydrogel scaffold and PLGA microspheres for supporting survival of neural stem cells. *Pharm Res* 2011; 28:1406–1414

This is the accepted but pre-publication version of the following article: Faulkner SD, Ruff CA, Fehlings MG. The potential for stem cells in cerebral palsy - piecing together the puzzle. *Semin Pediatr Neurol*. 2013 Jun;20(2):146-53. doi: 10.1016/j.spen.2013.06.002. Review. PubMed PMID: 23948689. which has been published in final format at: <http://www.sciencedirect.com/science/article/pii/S1071909113000284>

79. Wang L, Shansky J, Borselli C, et al.: Design and fabrication of a biodegradable, covalently crosslinked shape-memory alginate scaffold for cell and growth factor delivery. *Tissue Eng Part A* 2012; 18:2000–2007
80. Tysseling-Mattiace VM, Sahni V, Niece KL, et al.: Self-assembling nanofibers inhibit glial scar formation and promote axon elongation after spinal cord injury. *J Neurosci* 2008; 28:3814–3823[cited 2009 Dec 18]
81. Leung GKK, Wang YC, Wu W: Peptide nanofiber scaffold for brain tissue reconstruction. *Meth Enzymol* 2012; 508:177–190
82. Wright PA, Durham S, Ewins DJ, et al.: Neuromuscular electrical stimulation for children with cerebral palsy: a review. *Arch Dis Child* 2012; 97:364–371
83. Bolognini N, Vallar G, Casati C, et al.: Neurophysiological and behavioral effects of tDCS combined with constraint-induced movement therapy in poststroke patients. *Neurorehabil Neural Repair* 2011; 25:819–829